

Wightman function and Casimir densities on AdS bulk with application to the Randall–Sundrum braneworld

Aram A. Saharian*

*Department of Physics, Yerevan State University, 1 Alex Manoogian Str.,
375049 Yerevan, Armenia*

February 1, 2008

Abstract

Positive frequency Wightman function and vacuum expectation value of the energy-momentum tensor are computed for a massive scalar field with general curvature coupling parameter subject to Robin boundary conditions on two parallel plates located on $(D + 1)$ -dimensional AdS background. The general case of different Robin coefficients on separate plates is considered. The mode summation method is used with a combination of a variant of the generalized Abel-Plana formula for the series over zeros of combinations of cylinder functions. This allows us to extract manifestly the parts due to the AdS spacetime without boundaries and boundary induced parts. The asymptotic behavior of the vacuum densities near the plates and at large distances is investigated. The vacuum forces acting on the boundaries are presented as a sum of the self-action and interaction forces. The first one contains well-known surface divergences and needs further regularization. The interaction forces between the plates are attractive for Dirichlet scalar. We show that there is a region in the space of parameters defining the boundary conditions in which the interaction forces are repulsive for small distances and attractive for large distances. An application to the Randall-Sundrum braneworld with arbitrary mass terms on the branes is discussed.

1 Introduction

Anti-de Sitter (AdS) spacetime is one of the simplest and most interesting spacetimes allowed by general relativity. Quantum field theory in this background has been discussed by several authors (see, for instance, Refs. [1]–[17]). Much of early interest to AdS spacetime was motivated by the questions of principle related to the quantization of fields propagating on curved backgrounds. The importance of this theoretical work increased when it was realized that AdS spacetime emerges as a stable ground state solution in extended supergravity and Kaluza-Klein models and in string theories. The appearance of the AdS/CFT correspondence and braneworld models of Randall-Sundrum type has revived interest in this subject considerably. The AdS/CFT correspondence (for a review see [18]) represents a realization of the holographic principle and relates string theories or supergravity in the bulk of AdS with a conformal field theory living on its boundary. It has many interesting formal and physical facets and provides a powerful tool to investigate gauge field theories, in particular QCD. Recently it has been suggested that the introduction of compactified extra spatial dimensions may provide a solution to the hierarchy problem between the gravitational and electroweak mass scales [19, 20, 21]. The main idea to resolve the large hierarchy is that the small coupling of four dimensional gravity is generated by the large physical volume of extra dimensions. These theories provide a novel setting for

*Email: saharayan@server.physdep.r.am

discussing phenomenological and cosmological issues related to extra dimensions. The model introduced by Randall and Sundrum is particularly attractive. Their background solution consists of two parallel flat branes, one with positive tension and another with negative tension embedded in a five dimensional AdS bulk [20]. The fifth coordinate is compactified on S^1/Z_2 , and the branes are on the two fixed points. It is assumed that all matter fields are confined on the branes and only the gravity propagates freely in the five dimensional bulk. In this model, the hierarchy problem is solved if the distance between the branes is about 40 times the AdS radius and we live on the negative tension brane. More recently, scenarios with additional bulk fields have been considered [22, 23, 24, 25, 26].

In the scenario presented in [20] the distance between the branes is associated with the vacuum expectation value of a massless scalar field, called the radion. This modulus field has zero potential and consequently the distance is not determined by the dynamics of the model. For this scenario to be relevant, it is necessary to find a mechanism for generating a potential to stabilize the distance between the branes. Classical stabilization forces due to the non-trivial background configurations of a scalar field along an extra dimension were first discussed by Gell-Mann and Zwiebach [27]. With the revived interest in extra dimensions and braneworlds, as modified version of this mechanism, which exploits a classical force due to a bulk scalar field with different interactions with the branes, received significant attention [28] (and references therein). Another possibility for the stabilization mechanism arises due to the vacuum force generated by the quantum fluctuations about a constant background of a bulk field. The braneworld corresponds to a manifold with dynamical boundaries and all fields which propagate in the bulk will give Casimir-type contributions to the vacuum energy (for the Casimir effect see Refs. [29]–[34]), and as a result to the vacuum forces acting on the branes. In dependence of the type of a field and boundary conditions imposed, these forces can either stabilize or destabilize the braneworld. In addition, the Casimir energy gives a contribution to both the brane and bulk cosmological constants and, hence, has to be taken into account in the self-consistent formulation of the braneworld dynamics. Motivated by these, the role of quantum effects in braneworld scenarios has received some recent attention. For a conformally coupled scalar this effect was initially studied in Ref. [35] in the context of M-theory, and subsequently in Refs. [36]–[49] for a background Randall–Sundrum geometry (for the related heat kernel expansions see Refs. [50, 51, 52]). The models with dS branes are considered as well [48, 53, 54, 55]. For a conformally coupled bulk scalar the cosmological backreaction of the Casimir energy is investigated in Refs. [35, 48, 53, 56, 57, 58].

Investigation of local physical characteristics in the Casimir effect, such as expectation value of the energy-momentum tensor, is of considerable interest. In addition to describing the physical structure of the quantum field at a given point, the energy-momentum tensor acts as the source in the Einstein equations and therefore plays an important role in modelling a self-consistent dynamics involving the gravitational field. In this paper we will study the vacuum expectation value of the energy-momentum tensor of a scalar field with arbitrary curvature coupling parameter obeying Robin boundary conditions on two parallel plates in $(D+1)$ -dimensional AdS spacetime. The general case is considered when the constants in the Robin boundary conditions are different for separate plates. Robin type conditions are an extension of Dirichlet and Neumann boundary conditions and appear in a variety of situations, including the considerations of vacuum effects for a confined charged scalar field in external fields [59], spinor and gauge field theories, quantum gravity and supergravity [60, 61]. Robin conditions can be made conformally invariant, while purely-Neumann conditions cannot. Thus, Robin-type conditions are needed when one deals with conformally invariant theories in the presence of boundaries and wishes to preserve this invariance. It is interesting to note that the quantum scalar field satisfying the Robin condition on the boundary of the cavity violates the Bekenstein’s entropy-to-energy bound near certain points in the space of the parameter defining the boundary condition [62]. The Robin boundary

conditions are an extension of those imposed on perfectly conducting boundaries and may, in some geometries, be useful for depicting the finite penetration of the field into the boundary with the 'skin-depth' parameter related to the Robin coefficient. Mixed boundary conditions naturally arise for scalar and fermion bulk fields in the Randall-Sundrum model [22, 43]. To obtain the expectation values of the energy-momentum tensor, we first construct the positive frequency Wightman function. The application of the generalized Abel-Plana formula to the corresponding mode sum allows us to extract manifestly the boundary-free AdS part. The expressions for the boundary induced vacuum expectation values of the energy-momentum tensor are obtained by applying on the subtracted part a certain second order differential operator and taking the coincidence limit. Note that the Wightman function is also important in consideration of the response of particle detectors at a given state of motion (see, for instance, [63]).

We have organized the paper as follows. In the next section we consider the vacuum in the region between two parallel plates. The corresponding positive frequency Wightman function is evaluated by using the generalized Abel-Plana summation formula for the series over zeros of a combination of cylinder functions. This allows us to present the boundary induced part in terms of integrals with exponential convergence for the points away the boundaries. In Section 3 we consider the vacuum expectation value of the energy-momentum tensor for the case of a single plate. Both regions on the right and on the left from the plate are investigated. Various limiting cases are discussed. Section 4 is devoted to the vacuum energy-momentum tensor for the geometry of two parallel plates. The corresponding vacuum expectation values are presented in the form of the sum of single plates and 'interference' parts. The latter is finite everywhere including the points on the boundaries. The interaction forces between the plates are discussed as well. In Section 5 we show that the vacuum expectation value for the energy-momentum tensor of a bulk scalar in the Randall-Sundrum braneworld scenario is obtained from our results as a special case. The last section contains a summary of the work.

2 Wightman function

Consider a scalar field $\varphi(x)$ on background of a $(D+1)$ -dimensional plane-symmetric spacetime with the line element

$$ds^2 = g_{ik} dx^i dx^k = e^{-2\sigma(y)} \eta_{\mu\nu} dx^\mu dx^\nu - dy^2, \quad (2.1)$$

and with $\eta_{\mu\nu} = \text{diag}(1, -1, \dots, -1)$ being the metric for the D -dimensional Minkowski spacetime. Here and below $i, k = 0, 1, \dots, D$, and $\mu, \nu = 0, 1, \dots, D-1$. The corresponding field equation has the form

$$\left(g^{ik} \nabla_i \nabla_k + m^2 + \zeta R \right) \varphi(x) = 0, \quad (2.2)$$

where the symbol ∇_i is the operator for the covariant derivative associated with the metric g_{ik} , R is the corresponding Ricci scalar, and ζ is the curvature coupling parameter. For minimally and conformally coupled scalars one has $\zeta = 0$ and $\zeta = \zeta_c = (D-1)/4D$ correspondingly. Note that by making a coordinate transformation

$$z = \int e^{\sigma(y)} dy, \quad (2.3)$$

metric (2.1) is written in a conformally-flat form $ds^2 = e^{-2\sigma} \eta_{ik} dx^i dx^k$.

Below we will study quantum vacuum effects brought about by the presence of parallel infinite plane boundaries, located at $y = a$ and $y = b$, $a < b$, with mixed boundary conditions

$$\left(\tilde{A}_y + \tilde{B}_y \partial_y \right) \varphi(x) = 0, \quad y = a, b, \quad (2.4)$$

and constant coefficients $\tilde{A}_y, \tilde{B}_y$. The presence of boundaries modifies the spectrum for the zero-point fluctuations of the scalar field under consideration. This leads to the modification of the vacuum expectation values (VEVs) of physical quantities to compared with the case without boundaries. In particular, vacuum forces arise acting on the boundaries. This is well known Casimir effect. As a first stage, in this section we will consider the positive frequency Wightman function defined as the expectation value

$$G^+(x, x') = \langle 0 | \varphi(x) \varphi(x') | 0 \rangle, \quad (2.5)$$

where $|0\rangle$ is the amplitude for the vacuum state. In the next section we use this function to evaluate the VEVs for the energy-momentum tensor. Note that the Wightman function also determines the response of particle detectors in a given state of motion. By expanding the field operator over eigenfunctions and using the commutation relations one can see that

$$G^+(x, x') = \sum_{\alpha} \varphi_{\alpha}(x) \varphi_{\alpha}^*(x'), \quad (2.6)$$

where α denotes a set of quantum numbers, and $\{\varphi_{\alpha}(x)\}$ is a complete set of solutions to the field equation (2.2) satisfying boundary conditions (2.4).

On the base of the plane symmetry of the problem under consideration the corresponding eigenfunctions can be presented in the form

$$\varphi_{\alpha}(x^i) = \phi_{\mathbf{k}}(x^{\mu}) f_n(y), \quad (2.7)$$

where $\phi_{\mathbf{k}}(x^{\mu})$ are the standard Minkowskian modes on the hyperplane parallel to the plates:

$$\phi_{\mathbf{k}}(x^{\mu}) = \frac{e^{-i\eta_{\mu\nu} k^{\mu} x^{\nu}}}{\sqrt{2\omega(2\pi)^{D-1}}}, \quad k^{\mu} = (\omega, \mathbf{k}), \quad \omega = \sqrt{k^2 + m_n^2}, \quad k = |\mathbf{k}|. \quad (2.8)$$

Here the separation constants m_n are determined by boundary conditions (2.4) and will be given below. Substituting eigenfunctions (2.7) into the field equation (2.2) for the function $f_n(y)$ one obtains the following equation

$$-e^{D\sigma} \frac{d}{dy} \left(e^{-D\sigma} \frac{df_n}{dy} \right) + (m^2 + \zeta R) f_n = m_n^2 e^{2\sigma} f_n. \quad (2.9)$$

For the AdS geometry one has $\sigma(y) = k_D y$, $z = e^{\sigma(y)}/k_D$, and $R = -D(D+1)k_D^2$, where the AdS curvature radius is given by $1/k_D$. In this case the solution to equation (2.9) for the region $a < y < b$ is

$$f_n(y) = c_n e^{D\sigma/2} [J_{\nu}(m_n z) + b_{\nu} Y_{\nu}(m_n z)], \quad (2.10)$$

where $J_{\nu}(x)$, $Y_{\nu}(x)$ are the Bessel and Neumann functions, and

$$\nu = \sqrt{(D/2)^2 - D(D+1)\zeta + m^2/k_D^2}. \quad (2.11)$$

Here we will assume values of the curvature coupling parameter for which ν is real. For imaginary ν the ground state becomes unstable [4]. Note that for a conformally coupled massless scalar one has $\nu = 1/2$ and the cylinder functions in Eq. (2.10) are expressed via the elementary functions. From the boundary condition on $y = a$ one finds

$$b_{\nu} = -\frac{\bar{J}_{\nu}^{(a)}(m_n z_a)}{\bar{Y}_{\nu}^{(a)}(m_n z_a)}, \quad z_j = e^{\sigma(j)}/k_D, \quad j = a, b, \quad (2.12)$$

where we use the notation

$$\bar{F}^{(j)}(x) = A_j F(x) + B_j x F'(x), \quad A_j = \tilde{A}_j + \tilde{B}_j k_D D/2, \quad B_j = \tilde{B}_j k_D, \quad j = a, b \quad (2.13)$$

for a given function $F(x)$. Note that for Neumann scalar one has $A_j = B_j D/2$ and

$$\bar{F}^{(j)}(x) = F^{(N)}(x) \equiv x^{1-D/2} [x^{D/2} F(x)]', \quad \tilde{A}_j = 0. \quad (2.14)$$

From the boundary condition on the plane $y = b$ we receive that the eigenvalues m_n have to be solutions to the equation

$$C_\nu^{ab}(z_b/z_a, m_n z_a) \equiv \bar{J}_\nu^{(a)}(m_n z_a) \bar{Y}_\nu^{(b)}(m_n z_b) - \bar{Y}_\nu^{(a)}(m_n z_a) \bar{J}_\nu^{(b)}(m_n z_b) = 0. \quad (2.15)$$

We denote by $z = \gamma_{\nu,n}$, $n = 1, 2, \dots$, the zeros of the function $C_\nu^{ab}(\eta, z)$ in the right half-plane of the complex variable z , arranged in the ascending order, $\gamma_{\nu,n} < \gamma_{\nu,n+1}$. The eigenvalues for m_n are related to these zeros as

$$m_n = k_D \gamma_{\nu,n} e^{-\sigma(a)} = \gamma_{\nu,n} / z_a. \quad (2.16)$$

The coefficient c_n in Eq. (2.10) is determined from the orthonormality condition

$$\int_a^b dy e^{(2-D)\sigma} f_n(y) f_{n'}(y) = \delta_{nn'} \quad (2.17)$$

and is equal to

$$c_n^2 = \frac{\pi^2 u}{2k_D z_a^2} \bar{Y}_\nu^{(a)2}(u) T_\nu^{ab}(\eta, u), \quad u = \gamma_{\nu,n}, \quad \eta = z_b/z_a, \quad (2.18)$$

where we have introduced the notation

$$T_\nu^{ab}(\eta, u) = u \left\{ \frac{\bar{J}_\nu^{(a)2}(u)}{\bar{J}_\nu^{(b)2}(\eta u)} [A_b^2 + B_b^2(\eta^2 u^2 - \nu^2)] - A_a^2 + B_a^2(u^2 - \nu^2) \right\}^{-1}. \quad (2.19)$$

Note that, as we consider the quantization in the region between the branes, $z_a \leq z \leq z_b$, the modes defined by (2.10) are normalizable for all real values of ν from Eq. (2.11).

Substituting the eigenfunctions (2.7) into the mode sum (2.6), for the expectation value of the field product one finds

$$\langle 0 | \varphi(x) \varphi(x') | 0 \rangle = \frac{k_D^{D-1} (zz')^{D/2}}{2^{D+1} \pi^{D-3} z_a^2} \int d\mathbf{k} e^{i\mathbf{k}(\mathbf{x}-\mathbf{x}')} \sum_{n=1}^{\infty} h_\nu(\gamma_{\nu,n}) T_\nu^{ab}(\eta, \gamma_{\nu,n}), \quad (2.20)$$

where $\mathbf{x} = (x^1, x^2, \dots, x^{D-1})$ represents the coordinates in $(D-1)$ -hyperplane parallel to the plates and

$$h_\nu(u) = \frac{u e^{i\sqrt{u^2/z_a^2 + k^2}(t'-t)}}{\sqrt{u^2/z_a^2 + k^2}} g_\nu(u, uz/z_a) g_\nu(u, uz'/z_a). \quad (2.21)$$

Here and below we use the notation

$$g_\nu(u, v) = J_\nu(v) \bar{Y}_\nu^{(a)}(u) - \bar{J}_\nu^{(a)}(u) Y_\nu(v). \quad (2.22)$$

To sum over n we will use the summation formula derived in Refs. [64, 65] by making use of the generalized Abel-Plana formula [66, 64]. For a function $h(u)$ analytic in the right half-plane $\text{Re } u > 0$ this formula has the form

$$\begin{aligned} \frac{\pi^2}{2} \sum_{n=1}^{\infty} h(\gamma_{\nu,n}) T_\nu^{ab}(\eta, \gamma_{\nu,n}) &= \int_0^\infty \frac{h(x) dx}{\bar{J}_\nu^{(a)2}(x) + \bar{Y}_\nu^{(a)2}(x)} - \frac{\pi}{2} \text{Res}_{u=0} \left[\frac{h(u) \bar{H}_\nu^{(1b)}(\eta u)}{C_\nu^{ab}(\eta, u) \bar{H}_\nu^{(1a)}(u)} \right] - \\ &\quad - \frac{\pi}{4} \int_0^\infty \frac{\bar{K}_\nu^{(b)}(\eta x)}{\bar{K}_\nu^{(a)}(x) \bar{K}_\nu^{(a)}(x) \bar{I}_\nu^{(b)}(\eta x) - \bar{K}_\nu^{(b)}(\eta x) \bar{I}_\nu^{(a)}(x)} dx, \end{aligned} \quad (2.23)$$

where $I_\nu(u)$ and $K_\nu(u)$ are the Bessel modified functions. Formula (2.23) is valid for functions $h(u)$ satisfying the conditions

$$|h(u)| < \varepsilon_1(x)e^{c_1|y|} \quad |u| \rightarrow \infty, \quad u = x + iy, \quad (2.24)$$

and

$$h(ue^{\pi i}) = -h(u) + o(u^{-1}), \quad u \rightarrow 0, \quad (2.25)$$

where $c_1 < 2(\eta - 1)$, $x^{2\delta_{Ba^0-1}}\varepsilon_1(x) \rightarrow 0$ for $x \rightarrow +\infty$. Using the asymptotic formulae for the Bessel functions for large arguments when ν is fixed (see, e.g., [67]), we can see that for the function $h_\nu(u)$ from Eq. (2.21) the condition (2.24) is satisfied if $z + z' + |t - t'| < 2z_b$. In particular, this is the case in the coincidence limit $t = t'$ for the region under consideration, $z_a < z, z' < z_b$. As for $|u| < k$ one has $h_\nu(ue^{\pi i}) = -h_\nu(u)$, the condition (2.25) is also satisfied for the function $h_\nu(u)$. Note that $h_\nu(u) \sim u^{1-\delta_{k^0}}$ for $u \rightarrow 0$ and the residue term on the right of formula (2.23) vanishes. Applying to the sum over n in Eq. (2.20) formula (2.23), one obtains

$$\begin{aligned} \langle 0|\varphi(x)\varphi(x')|0\rangle &= \frac{k_D^{D-1}(zz')^{D/2}}{2^D\pi^{D-1}} \int d\mathbf{k} e^{i\mathbf{k}(\mathbf{x}-\mathbf{x}')} \left\{ \frac{1}{z_a^2} \int_0^\infty \frac{h_\nu(u)du}{\bar{J}_\nu^{(a)2}(u) + \bar{Y}_\nu^{(a)2}(u)} - \right. \\ &\quad \left. - \frac{2}{\pi} \int_k^\infty du u \frac{\Omega_{a\nu}(uz_a, uz_b)}{\sqrt{u^2 - k^2}} G_\nu^{(a)}(uz_a, uz) G_\nu^{(a)}(uz_a, uz') \cosh \left[\sqrt{u^2 - k^2}(t - t') \right] \right\}, \end{aligned} \quad (2.26)$$

where we have introduced notations

$$G_\nu^{(j)}(u, v) = I_\nu(v)\bar{K}_\nu^{(j)}(u) - \bar{I}_\nu^{(j)}(u)K_\nu(v), \quad j = a, b, \quad (2.27)$$

$$\Omega_{a\nu}(u, v) = \frac{\bar{K}_\nu^{(b)}(v)/\bar{K}_\nu^{(a)}(u)}{\bar{K}_\nu^{(a)}(u)\bar{I}_\nu^{(b)}(v) - \bar{K}_\nu^{(b)}(v)\bar{I}_\nu^{(a)}(u)} \quad (2.28)$$

(the function with $j = b$ will be used below). Note that we have assumed values of the coefficients A_α and B_α for which all zeros for Eq. (2.15) are real and have omitted the residue terms in the original formula in Refs. [64, 65]. In the following we will consider this case only.

To simplify the first term in the figure braces in Eq. (2.26), let us use the relation

$$\begin{aligned} \frac{g_\nu(u, uz/z_a)g_\nu(u, uz'/z_a)}{\bar{J}_\nu^{(a)2}(u) + \bar{Y}_\nu^{(a)2}(u)} &= J_\nu(uz/z_a)J_\nu(uz'/z_a) - \\ &- \frac{1}{2} \sum_{\beta=1}^2 \frac{\bar{J}_\nu^{(a)}(u)}{\bar{H}_\nu^{(\beta a)}(u)} H_\nu^{(\beta)}(uz/z_a) H_\nu^{(\beta)}(uz'/z_a) \end{aligned} \quad (2.29)$$

with $H_\nu^{(\beta)}(z)$, $\beta = 1, 2$, being the Hankel functions. Substituting this into the first integral in the figure braces of Eq. (2.26) we rotate the integration contour over u by the angle $\pi/2$ for $\beta = 1$ and by the angle $-\pi/2$ for $\beta = 2$. Under the condition $z + z' - |t - t'| > 2z_a$, the integrals over the arcs of the circle with large radius vanish. The integrals over $(0, ikz_a)$ and $(0, -ikz_a)$ cancel out and after introducing the Bessel modified functions one obtains

$$\begin{aligned} \int_0^\infty \frac{h_\nu(u)du}{\bar{J}_\nu^{(a)2}(u) + \bar{Y}_\nu^{(a)2}(u)} &= z_a^2 \int_0^\infty du u \frac{e^{i\sqrt{u^2 + k^2}(t'-t)}}{\sqrt{u^2 + k^2}} J_\nu(uz) J_\nu(uz') \\ &- \frac{2z_a^2}{\pi} \int_k^\infty du u \frac{\bar{I}_\nu^{(a)}(uz_a)}{\bar{K}_\nu^{(a)}(uz_a)} \frac{K_\nu(uz)K_\nu(uz')}{\sqrt{u^2 - k^2}} \cosh \left[\sqrt{u^2 - k^2}(t - t') \right]. \end{aligned} \quad (2.30)$$

Substituting this into formula (2.26), the Wightman function can be presented in the form

$$\begin{aligned} \langle 0|\varphi(x)\varphi(x')|0\rangle &= \langle 0_S|\varphi(x)\varphi(x')|0_S\rangle + \langle \varphi(x)\varphi(x') \rangle^{(a)} - \frac{k_D^{D-1}(zz')^{D/2}}{2^{D-1}\pi^D} \int d\mathbf{k} e^{i\mathbf{k}(\mathbf{x}-\mathbf{x}')} \\ &\times \int_k^\infty du u \frac{\Omega_{a\nu}(uz_a, uz_b)}{\sqrt{u^2 - k^2}} G_\nu^{(a)}(uz_a, uz) G_\nu^{(a)}(uz_a, uz') \cosh \left[\sqrt{u^2 - k^2}(t - t') \right]. \end{aligned} \quad (2.31)$$

Here the term

$$\langle 0_S | \varphi(x) \varphi(x') | 0_S \rangle = \frac{k_D^{D-1} (zz')^{D/2}}{2^D \pi^{D-1}} \int d\mathbf{k} e^{i\mathbf{k}(\mathbf{x}-\mathbf{x}')} \int_0^\infty du u \frac{e^{i\sqrt{u^2+k^2}(t'-t)}}{\sqrt{u^2+k^2}} J_\nu(uz) J_\nu(uz') \quad (2.32)$$

does not depend on the boundary conditions and is the Wightman function for the AdS space without boundaries. The second term on the right of Eq. (2.31),

$$\begin{aligned} \langle \varphi(x) \varphi(x') \rangle^{(a)} &= -\frac{k_D^{D-1} (zz')^{D/2}}{2^{D-1} \pi^D} \int d\mathbf{k} e^{i\mathbf{k}(\mathbf{x}-\mathbf{x}')} \times \\ &\times \int_k^\infty du u \frac{\bar{I}_\nu^{(a)}(uz_a)}{\bar{K}_\nu^{(a)}(uz_a)} \frac{K_\nu(uz) K_\nu(uz')}{\sqrt{u^2-k^2}} \cosh\left[\sqrt{u^2-k^2}(t-t')\right], \end{aligned} \quad (2.33)$$

does not depend on the parameters of the boundary at $z = z_b$ and is induced in the region $z > z_a$ by a single boundary at $z = z_a$ when the boundary $z = z_b$ is absent.

Note that expression (2.32) for the boundary-free Wightman function can also be written in the form

$$\langle 0_S | \varphi(x) \varphi(x') | 0_S \rangle = k_D^{D-1} (zz')^{D/2} \int_0^\infty dm m G_{M_D}^+(x^\mu, x'^\mu; m) J_\nu(mz) J_\nu(mz'), \quad (2.34)$$

where $G_{M_D}^+(x^\mu, x'^\mu; m)$ is the Wightman function for a scalar field with mass m in the D -dimensional Minkowski spacetime M_D . The right hand side in Eq. (2.34) can be further simplified by using the expression

$$G_{M_D}^+(x^\mu, x'^\mu; m) = \frac{m^{D/2-1}}{(2\pi)^{D/2}} \frac{K_{D/2-1}[m\sqrt{(\mathbf{x}-\mathbf{x}')^2 - (t-t'-i\varepsilon)^2}]}{[(\mathbf{x}-\mathbf{x}')^2 - (t-t'-i\varepsilon)^2]^{(D-2)/4}}, \quad (2.35)$$

with $\varepsilon > 0$. Substituting this into formula (2.34) and making use of the integration formula from [68], one obtains

$$\begin{aligned} \langle 0_S | \varphi(x) \varphi(x') | 0_S \rangle &= \frac{k_D^{D-1}}{(2\pi)^{(D+1)/2}} e^{-(D-1)\pi i/2} (v^2 - 1)^{(1-D)/4} Q_{\nu-1/2}^{(D-1)/2}(v) \\ &= \frac{k_D^{D-1} \Gamma(\nu + D/2) v^{-\nu-D/2}}{2^{D/2+\nu+1} \pi^{D/2} \Gamma(\nu + 1)} {}_2F_1\left(\frac{D+2\nu+2}{4}, \frac{D+2\nu}{4}; \nu+1; \frac{1}{v^2}\right), \end{aligned} \quad (2.36)$$

where $Q_\mu^\alpha(v)$ is the associated Legendre function of the second kind, ${}_2F_1(a, b; c; u)$ is the hypergeometric function (see, for instance, [67]), and

$$v = 1 + \frac{(z - z')^2 + (\mathbf{x} - \mathbf{x}')^2 - (t - t' - i\varepsilon)^2}{2zz'}. \quad (2.37)$$

It can be checked that in the limit $k_D \rightarrow 0$ from (2.32) the expression for the $(D+1)$ -dimensional Minkowski Wightman function is obtained. To see this note that in this limit $\nu \sim m/k_D \rightarrow \infty$, $k_D z \approx 1 + k_D y$ and, hence, as it follows from (2.37), one has $v \approx 1 + k_D^2 w^2/2$, where $w^2 = (y - y')^2 + (\mathbf{x} - \mathbf{x}')^2 - (t - t' - i\varepsilon)^2$. Using the integral representations for the functions $Q_\mu^\alpha(v)$ and $K_\mu(v)$ (formulae 8.8.2 and 9.6.23 in Ref. [67]), it can be seen that

$$\lim_{k_D \rightarrow 0} \left[k_D^{2\alpha} e^{-\alpha\pi i} (v^2 - 1)^{-\alpha/2} Q_{\nu-1/2}^\alpha(v) \right] = \frac{m^\alpha}{w^\alpha} K_\alpha(mw). \quad (2.38)$$

For $\alpha = (D-1)/2$ the expression on the right of this formula coincides with the $(D+1)$ -dimensional Minkowskian Wightman function up to the coefficient $(2\pi)^{-(D+1)/2}$. Hence, in

combination with the first formula in (2.36), relation (2.38) proves our statement. Alternatively, we can use formula (2.32), replacing the Bessel functions by their asymptotic expressions for large values of the order.

In the coincidence limit $x = x'$ expression (2.36) is finite for $D < 1$ and for the VEV of the field square one obtains

$$\langle 0_S | \varphi^2(x) | 0_S \rangle = \frac{k_D^{D-1}}{(4\pi)^{(D+1)/2}} \frac{\Gamma(\frac{1-D}{2}) \Gamma(D/2 + \nu)}{\Gamma(1 + \nu - D/2)}. \quad (2.39)$$

This quantity is independent on the spacetime point, which is a direct consequence of the maximal symmetry of the AdS bulk. Formula (2.39) coincides with the result of Refs. [10, 14] (a typo of [10] is corrected in [14]) obtained from the Feynman propagator in the coincidence limit (for the zeta function based calculations see [12, 16]). The expression on the right of Eq. (2.39) is analytic in the complex D -plane apart from simple poles coming from the gamma function in the nominator. Hence, it can be extended throughout the whole complex plane. For even D this expression is finite and according to the dimensional regularization procedure [63] can be taken as the regularized value for the field square. Formula (2.39) can be also obtained directly from (2.32) firstly integrating over $\mathbf{k}$ and by making use of the integration formula

$$\mathcal{I}_\nu(D) \equiv \int_0^\infty dx x^{D-1} J_\nu^2(x) = \frac{\Gamma(\frac{1-D}{2}) \Gamma(D/2 + \nu)}{2\sqrt{\pi} \Gamma(1 - D/2) \Gamma(1 + \nu - D/2)}. \quad (2.40)$$

By using the identity

$$\begin{aligned} \frac{\bar{I}_\nu^{(a)}(uz_a)}{\bar{K}_\nu^{(a)}(uz_a)} K_\nu(uz) K_\nu(uz') + \Omega_{a\nu}(uz_a, uz_b) G_\nu^{(a)}(uz_a, uz) G_\nu^{(a)}(uz_a, uz') = \\ \frac{\bar{K}_\nu^{(b)}(uz_b)}{\bar{I}_\nu^{(b)}(uz_b)} I_\nu(uz) I_\nu(uz') + \Omega_{b\nu}(uz_a, uz_b) G_\nu^{(b)}(uz_b, uz) G_\nu^{(b)}(uz_b, uz'), \end{aligned} \quad (2.41)$$

with

$$\Omega_{b\nu}(u, v) = \frac{\bar{I}_\nu^{(a)}(u) / \bar{I}_\nu^{(b)}(v)}{\bar{K}_\nu^{(a)}(u) \bar{I}_\nu^{(b)}(v) - \bar{K}_\nu^{(b)}(v) \bar{I}_\nu^{(a)}(u)}, \quad (2.42)$$

it can be seen that the Wightman function in the region $z_a \leq z \leq z_b$ can also be presented in the equivalent form

$$\begin{aligned} \langle 0 | \varphi(x) \varphi(x') | 0 \rangle &= \langle 0_S | \varphi(x) \varphi(x') | 0_S \rangle + \langle \varphi(x) \varphi(x') \rangle^{(b)} - \frac{k_D^{D-1} (zz')^{D/2}}{2^{D-1} \pi^D} \int d\mathbf{k} e^{i\mathbf{k}(\mathbf{x}-\mathbf{x}')} \\ &\times \int_k^\infty du u \frac{\Omega_{b\nu}(uz_a, uz_b)}{\sqrt{u^2 - k^2}} G_\nu^{(b)}(uz_b, uz) G_\nu^{(b)}(uz_b, uz') \cosh \left[\sqrt{u^2 - k^2} (t - t') \right], \end{aligned} \quad (2.43)$$

where

$$\begin{aligned} \langle \varphi(x) \varphi(x') \rangle^{(b)} &= - \frac{k_D^{D-1} (zz')^{D/2}}{2^{D-1} \pi^D} \int d\mathbf{k} e^{i\mathbf{k}(\mathbf{x}-\mathbf{x}')} \\ &\times \int_k^\infty du u \frac{\bar{K}_\nu^{(b)}(uz_b)}{\bar{I}_\nu^{(b)}(uz_b)} \frac{I_\nu(uz) I_\nu(uz')}{\sqrt{u^2 - k^2}} \cosh \left[\sqrt{u^2 - k^2} (t - t') \right] \end{aligned} \quad (2.44)$$

is the boundary part induced in the region $z < z_b$ by a single plate at $z = z_b$ when the plate $z = z_a$ is absent. Note that in the formulae given above the integration over angular part can be done by using the formula

$$\int d\mathbf{k} e^{i\mathbf{k}(\mathbf{x}-\mathbf{x}')} F(k) = (2\pi)^{(D-1)/2} \int_0^\infty dk k^{D-2} F(k) \frac{J_{(D-3)/2}(k|\mathbf{x}-\mathbf{x}'|)}{(k|\mathbf{x}-\mathbf{x}'|)^{(D-3)/2}}, \quad (2.45)$$

for a given function $F(k)$. Combining two forms, formulae (2.31) and (2.43), we see that the expressions for the Wightman function in the region $z_a \leq z \leq z_b$ is symmetric under the interchange $a \rightleftharpoons b$ and $I_\nu \rightleftharpoons K_\nu$. Note that the expression for the Wightman function is not symmetric with respect to the interchange of the plate indices. The reason for this is that, though the background AdS spacetime is homogeneous, the boundaries have nonzero extrinsic curvature tensors and two sides of the boundaries are not equivalent. In particular, for the geometry of a single brane the VEVs differ for the regions on the left and on the right of the brane. Here the situation is similar to that for the case of a spherical shell on background of the Minkowski spacetime.

3 Casimir densities for a single plate

In this section we will consider the VEV of the energy-momentum tensor for a scalar field in the case of a single plate located at $z = z_a$. As it has been shown in the previous section the Wightman function for this geometry is presented in the form

$$\langle 0|\varphi(x)\varphi(x')|0\rangle = \langle 0_S|\varphi(x)\varphi(x')|0_S\rangle + \langle \varphi(x)\varphi(x')\rangle^{(a)}, \quad (3.1)$$

where $|0\rangle$ is the amplitude for the corresponding vacuum state. The boundary induced part $\langle \varphi(x)\varphi(x')\rangle^{(a)}$ is given by formula (2.33) in the region $z > z_a$ and by formula (2.44) with replacement $z_b \rightarrow z_a$ in the region $z < z_a$. For points away from the plate this part is finite in the coincidence limit and in the corresponding formulae for the Wightman function we can directly put $x = x'$. Introducing a new integration variable $v = \sqrt{u^2 - k^2}$, transforming to the polar coordinates in the plane (v, k) and integrating over angular part, the following formula can be derived

$$\int_0^\infty dk k^{D-2} \int_k^\infty du \frac{uf(u)}{\sqrt{u^2 - k^2}} = \frac{\sqrt{\pi}\Gamma(\frac{D-1}{2})}{2\Gamma(D/2)} \int_0^\infty du u^{D-1} f(u). \quad (3.2)$$

By using this formula and Eq. (2.33), the boundary induced VEV for the field square in the region $z > z_a$ is presented in the form

$$\langle \varphi^2(x) \rangle^{(a)} = -\frac{k_D^{D-1} z^D}{2^{D-1} \pi^{D/2} \Gamma(D/2)} \int_0^\infty du u^{D-1} \frac{\bar{I}_\nu^{(a)}(uz_a)}{\bar{K}_\nu^{(a)}(uz_a)} K_\nu^2(uz), \quad z > z_a. \quad (3.3)$$

The corresponding formula in the region $z < z_a$ is obtained from Eq. (2.44) by a similar way and differs from Eq. (3.3) by replacements $I_\nu \rightleftharpoons K_\nu$.

By using the field equation, the expression for the metric energy-momentum tensor of a scalar field can be presented in the form

$$T_{ik} = \partial_i \varphi \partial_k \varphi + \left[\left(\zeta - \frac{1}{4} \right) g_{ik} \nabla_l \nabla^l - \zeta \nabla_i \nabla_k - \zeta R_{ik} \right] \varphi^2. \quad (3.4)$$

By virtue of this, for the VEV of the energy-momentum tensor we have

$$\langle 0|T_{ik}(x)|0\rangle = \lim_{x' \rightarrow x} \partial_i \partial'_k \langle 0|\varphi(x)\varphi(x')|0\rangle + \left[\left(\zeta - \frac{1}{4} \right) g_{ik} \nabla_l \nabla^l - \zeta \nabla_i \nabla_k - \zeta R_{ik} \right] \langle 0|\varphi^2(x)|0\rangle. \quad (3.5)$$

This corresponds to the point-splitting regularization technique. VEV (3.5) can be evaluated by substituting expression (3.1) for the positive frequency Wightman function and VEV of the field square into Eq. (3.5). First of all we will consider the region $z > z_a$. The vacuum energy-momentum tensor is diagonal and can be presented in the form

$$\langle 0|T_i^k|0\rangle = \langle 0_S|T_i^k|0_S\rangle + \langle T_i^k \rangle^{(a)}, \quad (3.6)$$

where

$$\langle 0_S | T_i^k | 0_S \rangle = \frac{k_D^{D+1} z^D \delta_i^k}{2^{D-1} \pi^{(D-1)/2} \Gamma(\frac{D-1}{2})} \int_0^\infty dk k^{D-2} \int_0^\infty du u \frac{\tilde{f}^{(i)}[J_\nu(uz)]}{\sqrt{u^2 + k^2}} \quad (3.7)$$

is the VEV for the energy-momentum tensor in the AdS background without boundaries, and the term

$$\langle T_i^k \rangle^{(a)} = -\frac{k_D^{D+1} z^D \delta_i^k}{2^{D-2} \pi^{(D+1)/2} \Gamma(\frac{D-1}{2})} \int_0^\infty dk k^{D-2} \int_k^\infty du u \frac{\bar{I}_\nu^{(a)}(uz_a)}{\bar{K}_\nu^{(a)}(uz_a)} \frac{\tilde{F}^{(i)}[K_\nu(uz)]}{\sqrt{u^2 - k^2}}, \quad (3.8)$$

is induced by a single boundary at $z = z_a$. For a given function $g(v)$ the functions $\tilde{F}^{(i)}[g(v)]$ in formula (3.8) are defined as

$$\begin{aligned} \tilde{F}^{(0)}[g(v)] &= \left(\frac{1}{2} - 2\zeta\right) \left[v^2 g'^2(v) + \left(D + \frac{4\zeta}{4\zeta - 1}\right) v g(v) g'(v) + \right. \\ &\quad \left. + (v^2 + \nu^2) g^2(v) \right] + (z^2 k^2 - v^2) g^2(v) \end{aligned} \quad (3.9)$$

$$\tilde{F}^{(i)}[g(v)] = \tilde{F}^{(0)}[g(v)] + \left(v^2 - \frac{D z^2 k^2}{D-1} \right) g^2(v), \quad i = 1, \dots, D-1, \quad (3.10)$$

$$\begin{aligned} \tilde{F}^{(D)}[g(v)] &= -\frac{v^2}{2} g'^2(v) + \frac{D}{2} (4\zeta - 1) v g(v) g'(v) + \\ &\quad + \frac{1}{2} [v^2 + \nu^2 + 2\zeta D(D+1) - D^2/2] g^2(v), \end{aligned} \quad (3.11)$$

and the expressions for the functions $\tilde{f}^{(i)}[g(v)]$ are obtained from those for $\tilde{F}^{(i)}[g(v)]$ by the replacement $v \rightarrow iv$. Note that the boundary-induced part (3.8) is finite for the points away the brane and, hence, the renormalization procedure is needed for the boundary-free part only. The latter is well-investigated in literature.

Using formula (3.2), it can be seen that the contribution of the second term on the right of Eq. (3.10) to the boundary part of the VEVs vanishes. From Eq. (3.8) now one obtains

$$\langle T_i^k \rangle^{(a)} = -\frac{k_D^{D+1} z^D \delta_i^k}{2^{D-1} \pi^{D/2} \Gamma(D/2)} \int_0^\infty du u^{D-1} \frac{\bar{I}_\nu^{(a)}(uz_a)}{\bar{K}_\nu^{(a)}(uz_a)} F^{(i)}[K_\nu(uz)], \quad z > z_a, \quad (3.12)$$

where the expressions for the functions $F^{(i)}[g(v)]$ directly follow from Eqs. (3.9)–(3.11) after the integration using formula (3.2):

$$\begin{aligned} F^{(i)}[g(v)] &= \left(\frac{1}{2} - 2\zeta\right) \left[v^2 g'^2(v) + \left(D + \frac{4\zeta}{4\zeta - 1}\right) v g(v) g'(v) + \right. \\ &\quad \left. + \left(\nu^2 + v^2 + \frac{2v^2}{D(4\zeta - 1)}\right) g^2(v) \right], \quad i = 0, 1, \dots, D-1, \end{aligned} \quad (3.13)$$

$$F^{(D)}[g(v)] = \tilde{F}^{(D)}[g(v)]. \quad (3.14)$$

By a similar way for the VEVs induced by a single brane in the region $z < z_a$, by making use of expression (2.44) (with replacement $z_b \rightarrow z_a$), one obtains

$$\langle T_i^k \rangle^{(a)} = -\frac{k_D^{D+1} z^D \delta_i^k}{2^{D-1} \pi^{D/2} \Gamma(D/2)} \int_0^\infty du u^{D-1} \frac{\bar{K}_\nu^{(a)}(uz_a)}{\bar{I}_\nu^{(a)}(uz_a)} F^{(i)}[I_\nu(uz)], \quad z < z_a. \quad (3.15)$$

Note that VEVs (3.12), (3.15) depend only on the ratio z/z_a which is related to the proper distance from the plate by equation

$$z/z_a = e^{k_D(y-a)}. \quad (3.16)$$

As we see, for the part of the energy-momentum tensor corresponding to the coordinates in the hyperplane parallel to the plates one has $\langle T_{\mu\nu} \rangle^{(a)} \sim \eta_{\mu\nu}$. Of course, we could expect this result from the problem symmetry. It can be seen that the VEVs obtained above obey the continuity equation $T_{i;k}^k = 0$ which for the AdS metric takes the form

$$z^{D+1} \frac{\partial}{\partial z} (z^{-D} T_D^D) + D T_0^0 = 0. \quad (3.17)$$

In the AdS part without boundary the integration over k can be done using the formula

$$\int_0^\infty \frac{k^{D-2} dk}{\sqrt{u^2 + k^2}} = \frac{u^{D-2}}{2\sqrt{\pi}} \Gamma\left(\frac{D-1}{2}\right) \Gamma\left(1 - \frac{D}{2}\right). \quad (3.18)$$

As for the boundary terms, the contribution corresponding to the second term on the right of Eq. (3.10) vanishes and one obtains

$$\langle 0_S | T_i^k | 0_S \rangle = \frac{k_D^{D+1} z^D \delta_i^k}{2^D \pi^{D/2}} \Gamma\left(1 - \frac{D}{2}\right) \int_0^\infty du u^{D-1} f^{(i)}[J_\nu(uz)], \quad (3.19)$$

where the expressions for the functions $f^{(i)}[g(v)]$ are obtained from those for $F^{(i)}[g(v)]$, Eqs. (3.13), (3.14), by the replacement $v \rightarrow iv$. Now from Eq. (3.13) we have the following relation between the vacuum energy density and pressures in the parallel directions (no summation over i)

$$\langle 0 | T_0^0 | 0 \rangle = \langle 0 | T_i^i | 0 \rangle, \quad i = 1, \dots, D-1. \quad (3.20)$$

To evaluate the u -integral in (3.19) we use formula (2.40) and the relations

$$\mathcal{I}_\nu(D+2) = \left(\nu^2 - \frac{D^2}{4}\right) \frac{D \mathcal{I}_\nu(D)}{D+1}, \quad \mathcal{I}_{\nu-1}(D+2) = \left(\nu - 1 - \frac{D}{2}\right) \left(\nu - \frac{D}{2}\right) \frac{D \mathcal{I}_\nu(D)}{D+1}, \quad (3.21)$$

$$\int_0^\infty dx x^D J_\nu(x) J_{\nu-1}(x) = \left(\nu - \frac{D}{2}\right) \mathcal{I}_\nu(D). \quad (3.22)$$

This yields

$$\int_0^\infty dx x^{D-1} f^{(i)}[J_\nu(x)] = \frac{m^2 \mathcal{I}_\nu(D)}{k_D^2 (D+1)}, \quad i = 0, 1, \dots, D. \quad (3.23)$$

As a result for the boundary-free AdS VEVs one receives

$$\langle 0_S | T_i^k | 0_S \rangle = \delta_i^k \frac{k_D^{D-1} m^2}{(4\pi)^{(D+1)/2}} \frac{\Gamma\left(\frac{1-D}{2}\right) \Gamma(D/2 + \nu)}{(D+1) \Gamma(1 + \nu - D/2)}. \quad (3.24)$$

Hence, the AdS VEVs are proportional to the corresponding metric tensor. Again, we could expect this result due to the maximally symmetry of the AdS background. Formula (3.24) can be obtained also directly from (2.39), by using the standard relation between the field square and the trace of the energy-momentum tensor.

For a conformally coupled massless scalar $\nu = 1/2$, and by making use of the expressions for the modified Bessel functions, it can be seen that $\langle T_i^k \rangle^{(a)} = 0$ in the region $z > z_a$ and

$$\langle T_D^D \rangle^{(a)} = -D \langle T_0^0 \rangle^{(a)} = -\frac{(k_D z/z_a)^{D+1}}{(4\pi)^{D/2} \Gamma(D/2)} \int_0^\infty \frac{t^D dt}{\frac{B_a(t-1)+2A_a}{B_a(t+1)-2A_a} e^t + 1} \quad (3.25)$$

in the region $z < z_a$. Note that the corresponding energy-momentum tensor for a single Robin plate in the Minkowski bulk vanishes [69] and the result for the region $z > z_a$ is obtained by a simple conformal transformation from that for the Minkowski case. In the region $z < z_a$ this

procedure does not work as in the AdS problem one has $0 < z < z_a$ instead of $-\infty < z < z_a$ in the Minkowski problem and, hence, the part of AdS under consideration is not a conformal image of the corresponding manifold in the Minkowski spacetime (for a general discussion of this question in conformally related problems see Ref. [70]).

The boundary induced VEVs given by equations (3.12) and (3.15), in general, can not be further simplified and need numerical calculations. Relatively simple analytic formulae for the brane-induced parts can be obtained in limiting cases. First of all, as a partial check, in the limit $k_D \rightarrow 0$ the corresponding formulae for a single plate on the Minkowski bulk are obtained (see Ref. [69] for the massless case and Ref. [71] for the massive case). This can be seen noting that in this limit $\nu \sim m/k_D$ is large and by introducing the new integration variable in accordance with $u = \nu y$, we can replace the Bessel modified functions by their uniform asymptotic expansions for large values of the order. The Minkowski result is obtained in the leading order.

In the limit $z \rightarrow z_a$ for a fixed k_D expressions (3.12) and (3.15) diverge. In accordance with (3.16), this corresponds to small proper distances from the brane. The surface divergences in the VEVs of the energy-momentum tensor are well-known in quantum field theory with boundaries and are investigated for various types of boundary geometries and boundary conditions (see, for instance, Refs. [72, 73]). Near the brane the main contribution into the integral over u in Eqs. (3.12), (3.15) comes from the large values of u and we can replace the Bessel modified functions by their asymptotic expressions for large values of the argument when the order is fixed (see, for instance, [67]). To the leading order this yields

$$\langle T_0^0 \rangle^{(a)} \sim \Gamma\left(\frac{D+1}{2}\right) \frac{Dk_D^{D+1}(\zeta - \zeta_c)\kappa(B_a)}{2^D \pi^{(D+1)/2} |1 - z_a/z|^{D+1}}, \quad (3.26)$$

$$\langle T_D^D \rangle^{(a)} \sim \langle T_0^0 \rangle^{(a)} \left(1 - \frac{z_a}{z}\right), \quad (3.27)$$

where we use the notation

$$\kappa(x) = \begin{cases} 1 & \text{if } x = 0, \\ -1 & \text{if } x \neq 0. \end{cases} \quad (3.28)$$

Note that the leading terms for the components with $i = 0, 1, \dots, D-1$ are symmetric with respect to the plate, and the $\frac{D}{2}$ - component has opposite signs for the different sides of the plate. Near the brane the vacuum energy densities have opposite signs for the cases of Dirichlet ($B_a = 0$) and non-Dirichlet ($B_a \neq 0$) boundary conditions. Recall that for a conformally coupled massless scalar the vacuum energy-momentum tensor vanishes in the region $z > z_a$ and is given by expression (3.25) in the region $z < z_a$. The latter is finite everywhere including the points on the plate.

For the points with the proper distances from the plate much larger compared with the AdS curvature radius one has $z \gg z_a$. This limit is important from the point of view of the application to the Randall-Sundrum braneworld. Introducing in Eq. (3.12) a new integration variable $y = zu$, by making use of formulae for the Bessel modified functions for small values of the argument, and assuming $A_a - B_a \nu \neq 0$, to the leading order we receive

$$\langle T_i^k \rangle^{(a)} \sim -\frac{k_D^{D+1} \delta_i^k 2^{2-D} \pi^{-D/2}}{\Gamma(D/2) \Gamma(\nu) \Gamma(\nu+1)} \frac{A_a + B_a \nu}{A_a - B_a \nu} \left(\frac{z_a}{2z}\right)^{2\nu} \int_0^\infty dy y^{D+2\nu-1} F^{(i)}[K_\nu(y)]. \quad (3.29)$$

The integral in this formula can be evaluated on the base of formulae

$$\int_0^\infty dx x^{\alpha-1} K_\nu^2(x) = \frac{2^{\alpha-3}}{\Gamma(\alpha)} \Gamma\left(\frac{\alpha}{2} + \nu\right) \Gamma\left(\frac{\alpha}{2} - \nu\right) \Gamma^2\left(\frac{\alpha}{2}\right), \quad (3.30)$$

$$\int_0^\infty dx x^{\alpha-1} K_\nu'^2(x) = \left[\frac{\alpha^2}{2} - \nu^2 - \frac{\alpha}{\alpha+1} \left(\frac{\alpha^2}{4} - \nu^2\right)\right] \int_0^\infty dx x^{\alpha-1} K_\nu^2(x). \quad (3.31)$$

This yields

$$\begin{aligned} \langle T_0^0 \rangle^{(a)} &\sim \frac{k_D^{D+1}}{(4\pi)^{(D-1)/2}} \frac{A_a + B_a \nu}{A_a - B_a \nu} \left(\frac{z_a}{2z} \right)^{2\nu} \left(4\zeta - \frac{D + 2\nu}{D + 2\nu + 1} \right) \\ &\times \frac{(2\nu - 1)\Gamma(D/2 + \nu + 1)\Gamma(D/2 + 2\nu)}{\Gamma(\nu + 1)\Gamma(D/2 + 1/2 + \nu)}, \end{aligned} \quad (3.32)$$

$$\langle T_D^D \rangle^{(a)} \sim \frac{D}{D + 2\nu} \langle T_0^0 \rangle^{(a)}, \quad z \gg z_a, \quad (3.33)$$

and the boundary-induced VEVs are exponentially suppressed by the factor $\exp[2\nu k_D(a - y)]$. In the limit under consideration the ratio of the energy density and the perpendicular pressure is a negative constant. We see that in the case $\zeta \leq \zeta_c$, at large distances from the plate the energy density is positive for $|B_a/A_a| > 1/\nu$ and is negative for $|B_a/A_a| < 1/\nu$. Combining this with the asymptotic behavior near the plate, Eq. (3.26), we conclude that for $B_a \neq 0$ and $|B_a/A_a| < 1/\nu$ the energy density is positive near the plate and is negative at large distances approaching to zero and, hence, for some value of z it has a minimum with the negative energy density.

Now we turn to the limit $z \ll z_a$ for a fixed k_D . This corresponds to the points near the AdS boundary presented by the hyperplane $z = 0$ with the proper distances from the brane much larger compared with the AdS curvature radius. Introducing in Eq. (3.15) a new integration variable $y = uz_a$ and by making use of the formulae for the Bessel modified functions for small values of the argument, to the leading order one receives

$$\begin{aligned} \langle T_0^0 \rangle^{(a)} &\sim -\frac{k_D^{D+1}}{\pi^{D/2}\Gamma(D/2)} \left(\frac{z}{2z_a} \right)^{D+2\nu} \frac{D + 2\nu - 4\zeta(D + 2\nu + 1)}{\Gamma(\nu)\Gamma(\nu + 1)} \\ &\times \int_0^\infty dx x^{D+2\nu-1} \frac{\bar{K}_\nu^{(a)}(x)}{\bar{I}_\nu^{(a)}(x)}, \end{aligned} \quad (3.34)$$

$$\langle T_D^D \rangle^{(a)} \sim -\frac{D}{2\nu} \langle T_0^0 \rangle^{(a)}, \quad z \ll z_a. \quad (3.35)$$

In this limit the exponential suppression takes place by the factor $\exp[(D + 2\nu)k_D(y - a)]$. The ratio of the energy density and the perpendicular pressure is a positive constant. Due to the well-known properties of the Bessel modified functions, the integral in Eq. (3.34) is positive for small values of the ratio B_a/A_a and is negative for large values and, hence, as in the previous case there is a possibility for the change of the sign for the energy density as a function on the distance from the boundary. Now by taking into account relation (3.16), from Eqs. (3.32), (3.34) we conclude that for large proper distances from the plate to compared with the AdS curvature radius the boundary induced parts are exponentially suppressed.

In the large mass limit, $m \gg k_D$, from (2.11) one has $\nu \approx m/k_D \gg 1$. To find the leading terms of the corresponding asymptotic expansions for the vacuum energy-momentum tensor components we replace the integration variable in Eqs. (3.12) and (3.15), $x = \nu t$, and use the uniform asymptotic expansions for the Bessel modified functions for large values of the order. The main contribution to the integrals over t comes from small values of t . For a fixed z to the leading order one receives

$$\langle T_0^0 \rangle^{(a)} \sim (4\zeta - 1)\kappa(B_a) \frac{k_D^{D+1}}{\pi^{D/2}} \left(\frac{m}{2k_D} \right)^{D/2+1} \frac{e^{-2|\ln(z_a/z)|m/k_D}}{|1 - z_a^2/z^2|^{D/2}}, \quad (3.36)$$

$$\langle T_D^D \rangle^{(a)} \sim \frac{Dk_D}{2m} \langle T_0^0 \rangle^{(a)}, \quad m \gg k_D, \quad (3.37)$$

and we have an exponential suppression of the vacuum expectation values for large values of the mass.

For large values of the parameter k_D and fixed proper distances from the plate, $k_D|y-a| \gg 1$, from relation (3.16) one has $z_a \ll z$ or $z_a \gg z$. Hence, in this limit, which corresponds to strong gravitational fields, we have an asymptotic behavior described by Eqs. (3.32), (3.33) in the region $z > z_a$ and by Eqs. (3.34), (3.35) in the region $z < z_a$ with the exponential suppression of the boundary induced parts by the factor $\exp[2\nu k_D(a-y)]$ in the first case and by the factor $\exp[(2\nu + D)k_D(y-a)]$ in the second case.

In figures 1 and 2 we have plotted the vacuum energy density [curves (a)] and $\frac{D}{D-1}$ – stress [curves (b)] induced by a single plate, as functions on $k_D(y-a)$ for a minimally coupled massless scalar field in $D = 3$. In figure 1 the cases of Dirichlet (left panel) and Neumann (right panel) boundary conditions are presented. The energy density is negative everywhere for Dirichlet scalar and is positive for Neumann scalar. It can be seen that for both Dirichlet and Neumann cases the boundary-induced energy-momentum tensors violate the strong energy condition. The weak energy condition is violated by Dirichlet scalar and is satisfied by Neumann scalar. The energy conditions play a key role in the singularity theorems of general relativity and these properties may have important consequences in the consideration of the gravitational back reaction of vacuum quantum effects. Figure 2 corresponds to the Robin boundary condition with $B_a/A_a = 1/4$ and illustrates the possibility for the change of the sign for the energy density as a function on distance from the plate: the energy density is positive near the plate and is negative for large distances from the plate. Note that the ratio $\langle T_D^D \rangle^{(a)} / \langle T_0^0 \rangle^{(a)}$ is a coordinate dependent function. In accordance with the asymptotic estimates given above, this ratio tends to zero for the points near the brane and to the constant values $D/(D+2\nu)$ and $-D/(2\nu)$ in the limits $z \gg z_a$ and $z \ll z_a$, respectively.




Figure 1: Boundary induced vacuum densities $\langle T_0^0 \rangle^{(a)} / k_D^{D+1}$ and $\langle T_D^D \rangle^{(a)} / k_D^{D+1}$ as functions on $k_D(y-a)$ for a minimally coupled massless scalar in $D = 3$. The curves (a) correspond to the energy density and the curves (b) correspond to the $\frac{D}{D-1}$ – component of the vacuum stress. The left panel is for Dirichlet scalar and the right one is for Neumann scalar.

4 Two-plates geometry

4.1 Vacuum densities

In this section we will investigate the VEVs for the energy-momentum tensor in the region between two parallel plates. Substituting the corresponding Wightman function from Eq. (2.31)




Figure 2: The same as in figure 1 for Robin scalar with $B_a/A_a = 1/4$.

into the mode sum formula (3.5), we obtain

$$\begin{aligned} \langle 0|T_i^k|0\rangle &= \langle 0_S|T_i^k|0_S\rangle + \langle T_i^k\rangle^{(a)} - \frac{k_D^{D+1}z^D\delta_i^k}{2^{D-2}\pi^{(D+1)/2}\Gamma\left(\frac{D-1}{2}\right)} \int_0^\infty dk k^{D-2} \\ &\times \int_k^\infty du u \frac{\Omega_{av}(uz_a, uz_b)}{\sqrt{u^2 - k^2}} \tilde{F}^{(i)}\left[G_\nu^{(a)}(uz_a, uz)\right], \quad z_a < z < z_b \end{aligned} \quad (4.1)$$

where for a given function $g(v)$ the functions $\tilde{F}^{(i)}[g(v)]$ are defined in accordance with Eqs. (3.9)–(3.11). By using formula (3.2), we see that the contribution of the second term on the right of Eq. (3.10) vanishes and from Eq. (4.1) one obtains

$$\langle 0|T_i^k|0\rangle = \langle 0_S|T_i^k|0_S\rangle + \langle T_i^k\rangle^{(a)} - \frac{k_D^{D+1}z^D\delta_i^k}{2^{D-1}\pi^{\frac{D}{2}}\Gamma\left(\frac{D}{2}\right)} \int_0^\infty du u^{D-1} \Omega_{av}(uz_a, uz_b) F^{(i)}\left[G_\nu^{(a)}(uz_a, uz)\right], \quad (4.2)$$

with the functions $F^{(i)}[g(v)]$ from Eqs. (3.13), (3.14).

By making use of the Wightman function in form (2.43), we obtain an alternative representation of the VEVs:

$$\langle 0|T_i^k|0\rangle = \langle 0_S|T_i^k|0_S\rangle + \langle T_i^k\rangle^{(b)} - \frac{k_D^{D+1}z^D\delta_i^k}{2^{D-1}\pi^{\frac{D}{2}}\Gamma\left(\frac{D}{2}\right)} \int_0^\infty du u^{D-1} \Omega_{bv}(uz_a, uz_b) F^{(i)}\left[G_\nu^{(b)}(uz_b, uz)\right]. \quad (4.3)$$

This form is obtained from (4.2) by the replacements $a \rightleftharpoons b$, $I_\nu \rightleftharpoons K_\nu$. The vacuum energy-momentum tensor in the region between the branes is not symmetric under the interchange of indices of the branes. As it has been mentioned above, the reason for this is that, though the background spacetime is homogeneous, due to the non-zero extrinsic curvature tensors for the branes, the regions on the left and on the right of the brane are not equivalent. By the same way as for the case of a single plate, it can be seen that in the limit $k_D \rightarrow 0$ from formulae (4.2) and (4.3) the corresponding results for two plates geometry in the Minkowski bulk (see Refs. [69, 71]) are obtained.

On the base of formula (4.2), the VEVs in the region $z_a < z < z_b$ can be written in the form

$$\langle 0|T_i^k|0\rangle = \langle 0_S|T_i^k|0_S\rangle + \langle T_i^k\rangle^{(a)} + \langle T_i^k\rangle^{(b)} + \Delta\langle T_i^k\rangle, \quad (4.4)$$

with the 'interference' term

$$\begin{aligned} \Delta \langle T_i^k \rangle &= -\frac{k_D^{D+1} z^D \delta_i^k}{2^{D-1} \pi^{D/2} \Gamma(D/2)} \int_0^\infty du u^{D-1} \times \\ &\times \left\{ \Omega_{av}(uz_a, uz_b) F^{(i)}[G_\nu^{(a)}(uz_a, uz)] - \frac{\bar{K}_\nu^{(b)}(uz_b)}{\bar{I}_\nu^{(b)}(uz_b)} F^{(i)}[I_\nu(uz)] \right\}. \end{aligned} \quad (4.5)$$

Another form we can obtain by using formulae (3.12) and (4.3). Note that the surface divergences are contained in the second and third terms on the right of Eq. (4.4) and expression (4.5) is finite for all values $z_a \leq z \leq z_b$ (for large values u the subintegrand behaves as $u^D e^{2u(z_a - z_b)}$). For the proper distances between the plates much less than the AdS curvature radius, $k_D(b - a) \ll 1$, one has $(1 - z_a/z_b) \ll 1$ and the main contribution into the integral in Eq. (4.5) comes from large values of u . Replacing the Bessel modified functions by their asymptotic expansions for large values of the argument, for the VEVs from Eq. (4.5) to the leading order one receives

$$\Delta \langle T_D^D \rangle \sim \frac{D \Gamma(\frac{D+1}{2}) \zeta_R(D+1)}{(4\pi)^{(D+1)/2} (b-a)^{D+1}} \frac{1 - 2^{-D-1} [1 - \kappa(B_a) \kappa(B_b)]}{\kappa(B_a) \kappa(B_b)} \quad (4.6)$$

$$\begin{aligned} \Delta \langle T_0^0 \rangle &\sim -\frac{1}{D} \Delta \langle T_D^D \rangle - \frac{(\zeta - \zeta_c)}{2^{2D-1} \pi^{D/2} \Gamma(D/2) (b-a)^{D+1}} \times \\ &\times \int_0^\infty \frac{t^D dt}{\kappa(B_a) \kappa(B_b) e^t - 1} \left[\kappa(B_a) \exp\left(t \frac{z_a - z}{z_b - z_a}\right) + \kappa(B_b) \exp\left(t \frac{z - z_b}{z_b - z_a}\right) \right] \end{aligned} \quad (4.7)$$

where $\zeta_R(z)$ is the Riemann zeta function and the function $\kappa(x)$ is defined by Eq. (3.28). The leading terms given by formulae (4.6) and (4.7) are the same as for the corresponding quantities on Minkowski spacetime background. In particular, the 'interference' part of the $\frac{D}{D} -$ component does not depend on the curvature coupling parameter. In the limit of large distances between the plates, when $k_D(b - a) \gg 1$, introducing a new integration variable $v = uz_b$ and expanding over z_a/z_b , to the leading order from Eq. (4.5) one obtains

$$\Delta \langle T_i^k \rangle \sim \delta_i^k \left(\frac{z_a}{z_b} \right)^{2\nu} g^{(i)} \left(\frac{z}{z_b} \right), \quad (4.8)$$

where the form of the function $g^{(i)}(v)$ can be found from Eq. (4.5). For $v \ll 1$ one has $g(v) \sim v^D$ and from (4.8) it follows that the 'interference' part of the vacuum energy-momentum tensor is exponentially suppressed for large interbrane distances.

4.2 Interaction forces

Now we turn to the interaction forces between the plates. The vacuum force acting per unit surface of the plate at $z = z_j$ is determined by the $\frac{D}{D} -$ component of the vacuum energy-momentum tensor at this point. The corresponding effective pressures can be presented as a sum of two terms:

$$p^{(j)} = p_1^{(j)} + p_{(\text{int})}^{(j)}, \quad j = a, b. \quad (4.9)$$

The first term on the right is the pressure for a single plate at $z = z_j$ when the second plate is absent. This term is divergent due to the surface divergences in the vacuum expectation values. The second term on the right of Eq. (4.9),

$$p_{(\text{int})}^{(j)} = -\langle T_D^D \rangle^{(j_1)} - \Delta \langle T_D^D \rangle, \quad z = z_j, \quad j, j_1 = a, b, \quad j_1 \neq j, \quad (4.10)$$

is the pressure induced by the presence of the second plate, and can be termed as an interaction force. It is determined by the last terms on the right of formulae (4.2) and (4.3) for the plate at $z = z_a$ and $z = z_b$ respectively. Substituting into these terms $z = z_j$ and using relations

$$G_\nu^{(j)}(u, u) = -B_j, \quad G_\nu^{(j)'}(u, u) = A_j/u, \quad (4.11)$$

one has

$$p_{(\text{int})}^{(j)} = \frac{k_D^{D+1}}{2^D \pi^{D/2} \Gamma(\frac{D}{2})} \int_0^\infty dx x^{D-1} \Omega_{j\nu}(xz_a/z_j, xz_b/z_j) \\ \times \{ (x^2 - \nu^2 + 2m^2/k_D^2) B_j^2 - D(4\zeta - 1) A_j B_j - A_j^2 \}. \quad (4.12)$$

Note that due to the asymmetry in the VEV of the energy-momentum tensor, the interaction forces acting on the branes are not symmetric under the interchange of the brane indices. By taking into account that $K_\nu(u)I_\nu(v) - K_\nu(v)I_\nu(u) > 0$ and $K'_\nu(u)I'_\nu(v) - K'_\nu(v)I'_\nu(u) < 0$ for $u < v$, it can be easily seen that the vacuum effective pressures are negative for Dirichlet scalar and for a scalar with $A_a = A_b = 0$ and, hence, the corresponding interaction forces are attractive for all values of the interplate distance. In figure 3 these forces are plotted as functions on z_a/z_b for massless minimally coupled Dirichlet and Neumann scalars. As it has been shown above, in the case of a conformally coupled massless scalar field for the first term on the right of Eq. (4.9) one has $p_1^{(a)} = 0$ and $p_1^{(b)}$ is determined from Eq. (3.25) with the replacement $a \rightarrow b$ (recall that we consider the region $z_a \leq z \leq z_b$). It can be seen that the corresponding formulae for $p^{(j)}$ obtained from Eqs. (4.9), (4.12) coincide with those given in Ref. [47]. Note that in this case $p^{(a)}/z_a^{D+1} = p^{(b)}/z_b^{D+1}$. Using the Wronskian for the Bessel modified functions, it can be seen that

$$[B_j^2(x^2 z_j^2 + \nu^2) - A_j^2] \Omega_{j\nu}(xz_a, xz_b) = n_j z_j \frac{\partial}{\partial z_j} \ln \left| 1 - \frac{\bar{I}_\nu^{(a)}(xz_a) \bar{K}_\nu^{(b)}(xz_b)}{\bar{I}_\nu^{(b)}(xz_b) \bar{K}_\nu^{(a)}(xz_a)} \right|, \quad j = a, b, \quad (4.13)$$

where $n_a = 1$, $n_b = -1$. This allows us to write the expressions (4.12) for the interaction forces per unit surface in another equivalent form:

$$p_{(\text{int})}^{(j)} = \frac{n_j k_D^{D+1} z_j^{D+1}}{2^D \pi^{D/2} \Gamma(\frac{D}{2})} \int_0^\infty dx x^{D-1} \left[1 + D B_j \frac{2\zeta B_j + (1 - 4\zeta) \tilde{A}_j}{B_j^2(x^2 z_j^2 + \nu^2) - A_j^2} \right] \\ \times \frac{\partial}{\partial z_j} \ln \left| 1 - \frac{\bar{I}_\nu^{(a)}(xz_a) \bar{K}_\nu^{(b)}(xz_b)}{\bar{I}_\nu^{(b)}(xz_b) \bar{K}_\nu^{(a)}(xz_a)} \right|. \quad (4.14)$$

Note that these forces in general are different for $j = a$ and $j = b$. For Dirichlet scalar the second term in the square brackets is zero.

Let us consider the limiting cases for the interaction forces described by Eq. (4.12). For small distances compared with the AdS curvature radius, $k_D(b - a) \ll 1$, the leading terms are the same as for the plates in the Minkowski bulk:

$$p_{(\text{int})}^{(j)} = D(1 - 2^{-D}) \frac{\Gamma(\frac{D+1}{2}) \zeta_R(D+1)}{(4\pi)^{(D+1)/2} (b-a)^{D+1}}, \quad (4.15)$$

in the case of Dirichlet boundary condition on one plate and non-Dirichlet boundary condition on the another, and

$$p_{(\text{int})}^{(j)} = -D \frac{\Gamma(\frac{D+1}{2}) \zeta_R(D+1)}{(4\pi)^{(D+1)/2} (b-a)^{D+1}}, \quad (4.16)$$




Figure 3: The vacuum effective pressures $(z_b/z_j)^D p_{(\text{int})}^{(j)}/k_D^{D+1}$, $j = a, b$, as functions on z_a/z_b for a massless minimally coupled scalar field in $D = 3$ determining the interaction forces between Dirichlet (left panel) and Neumann (right panel) plates. Curves a and b correspond to the pressures on the plates $z = z_j$, $j = a, b$ respectively.

for all other cases. Note that in the first case the interaction forces are repulsive and are attractive for the second case. For large distances between the plates, $k_D(b - a) \gg 1$ (this limit is realized in the Randall-Sundrum model), by using the expressions for the modified Bessel functions for small values of the argument, one finds

$$p_{(\text{int})}^{(a)} \sim \left(\frac{z_a}{2z_b}\right)^{D+2\nu} \frac{4k_D^{D+1} [(2m^2/k_D^2 - \nu^2)B_a^2 - D(4\zeta - 1)A_a B_a - A_a^2]}{\pi^{D/2}\Gamma(D/2)\Gamma^2(\nu)(B_a\nu - A_a)^2} \times \int_0^\infty dx x^{2\nu+D-1} \frac{\bar{K}_\nu^{(b)}(x)}{\bar{I}_\nu^{(b)}(x)}, \quad (4.17)$$

$$p_{(\text{int})}^{(b)} \sim \left(\frac{z_a}{2z_b}\right)^{2\nu} \frac{A_a + B_a\nu}{A_a - B_a\nu} \frac{k_D^{D+1}}{2^{D-1}\pi^{D/2}\Gamma(D/2)\nu\Gamma^2(\nu)} \times \int_0^\infty dx \frac{x^{2\nu+D-1}}{\bar{I}_\nu^{(b)2}(x)} [(x^2 - \nu^2 + 2m^2/k_D^2)B_b^2 - D(4\zeta - 1)A_b B_b - A_b^2]. \quad (4.18)$$

In dependence of the values for the coefficients in the boundary conditions, these effective pressures can be either positive or negative, leading to repulsive or attractive forces at large distances. To illustrate these possibilities, in figure 4 we have plotted the effective pressures as functions on z_a/z_b for Robin boundary conditions. In the case of the left panel, the vacuum interaction forces are attractive for small distances and repulsive for large distances. For the right panel we have an opposite situation, the interaction forces are attractive for large distances and repulsive for small distances. For the latter case the vacuum interaction forces provide a possibility for a stabilization of the interplate distance. The dependence of the vacuum effective pressures on the ratio z_a/z_b for various values of the Robin coefficient B_b is given in figures 5 and 6.

5 Application to the Randall-Sundrum braneworld

In this section we will consider the application of the results obtained in the previous sections to the Randall-Sundrum braneworld model [20] based on the AdS geometry with one extra




Figure 4: The same as in figure 3 for mixed boundary conditions. Left panel corresponds to $\tilde{A}_a = 0$, $B_a = 1$, $\tilde{A}_b = 1$, $B_b = 0.2$ and right panel corresponds to $\tilde{A}_a = 1$, $B_a = 0$, $\tilde{A}_b = 1$, $B_b = 0.2$.




Figure 5: The $D = 3$ vacuum effective pressures $(z_b/z_j)^D p_{\text{int}}^{(j)}/k_D^{D+1}$, $j = a, b$, for a massless minimally coupled scalar field as functions on z_a/z_b and B_b . The values for the other Robin coefficients are $\tilde{A}_a = 1$, $B_a = 0$, and $\tilde{A}_b = 1$. Left panel corresponds to $j = a$ and right panel corresponds to $j = b$.




Figure 6: The same as in figure 5 for the values of Robin coefficients $\tilde{A}_a = 0$, $B_a = 1$, and $\tilde{A}_b = 1$.

dimension. The fifth dimension y is compactified on an orbifold, S^1/Z_2 of length L , with $-L \leq y \leq L$. The orbifold fixed points at $y = 0$ and $y = L$ are the locations of two 3-branes. Below we will allow these submanifolds to have an arbitrary dimension D . The metric in the Randall-Sundrum model has the form (2.1) with

$$\sigma(y) = k_D |y|. \quad (5.1)$$

For the corresponding Ricci scalar one has

$$R = 4Dk_D[\delta(y) - \delta(y - L)] - D(D + 1)k_D^2. \quad (5.2)$$

Let us consider a bulk scalar φ with the curvature coupling parameter ζ and with the action functional

$$S = \frac{1}{2} \int d^D x dy \sqrt{|g|} \left\{ g^{ik} \partial_i \varphi \partial_k \varphi - [m^2 + c_1 \delta(y) + c_2 \delta(y - L) + \zeta R] \varphi^2 \right\}, \quad (5.3)$$

where m is the bulk mass, c_1 and c_2 are the brane mass terms on the branes $y = 0$ and $y = L$ respectively. The field equation obtained from (5.3) is

$$\nabla_i \nabla^i \varphi + [m^2 + c_1 \delta(y) + c_2 \delta(y - L) + \zeta R] \varphi = 0. \quad (5.4)$$

The corresponding eigenfunctions can be written in form (2.7) and (2.8), where $f_n(y)$ is a solution to the equation

$$\begin{aligned} -e^{D\sigma} \frac{d}{dy} \left(e^{-D\sigma} \frac{df_n}{dy} \right) + [m^2 + (c_1 + 4D\zeta k_D) \delta(y) + (c_2 - 4D\zeta k_D) \delta(y - L) - \\ - D(D + 1)\zeta k_D^2] f_n(y) = m_n^2 e^{2\sigma} f_n(y). \end{aligned} \quad (5.5)$$

To obtain the boundary conditions for the function $f_n(y)$ we integrate (5.5) first about $y = 0$ and then about $y = L$. Assuming that the function $f_n(y)$ is continuous at these points, we receive

$$\lim_{\epsilon \rightarrow 0} \frac{df_n(y)}{dy} \Big|_{y=y_j - (-1)^j \epsilon}^{y=y_j + (-1)^j \epsilon} = [c_j - (-1)^j 4D\zeta k_D] f_n(y_j), \quad j = 1, 2, \quad y_1 = 0, \quad y_2 = L, \quad (5.6)$$

with $\epsilon > 0$. First consider the case of the untwisted scalar field for which $f_n(y) = f_n(-y)$, and the solution to Eq. (5.5) for $y \neq 0, L$ is given by expression (2.10) with $z = e^{k_D|y|}$. The boundary conditions which follow from relations (5.6) are in form (2.4) with (see also Ref. [43] for the case $c_1 = c_2 = 0$)

$$\frac{\tilde{A}_a}{\tilde{B}_a} = -\frac{1}{2}(c_1 + 4D\zeta k_D), \quad \frac{\tilde{A}_b}{\tilde{B}_b} = -\frac{1}{2}(-c_2 + 4D\zeta k_D), \quad (5.7)$$

and respectively

$$\frac{A_a}{B_a} = \frac{1}{2}[D(1 - 4\zeta) - c_1/k_D], \quad \frac{A_b}{B_b} = \frac{1}{2}[D(1 - 4\zeta) + c_2/k_D]. \quad (5.8)$$

For the twisted scalar $f_n(-y) = -f_n(y)$ and from the Z_2 identification on S^1 one has $f_n(0) = f_n(L) = 0$ and, hence, $\tilde{B}_a = \tilde{B}_b = 0$, which correspond to Dirichlet boundary conditions on both branes. The normalization condition for the eigenfunctions $f_n(y)$ now has the form

$$\int_{-L}^L dy e^{(2-D)\sigma(y)} f_n(y) f_{n'}(y) = \delta_{nn'}. \quad (5.9)$$

As a result the normalization coefficient c_n differs from (2.15) by additional factor $1/2$. Hence, the Wightman function in the Randall-Sundrum braneworld is given by formulae (2.31) or equivalently by (2.43) with an additional factor $1/2$ and with Robin coefficients given by Eq. (5.7). Similarly, the VEV of the energy-momentum tensor is defined by formulae (4.2) or (4.3) by an additional factor $1/2$. Note that global quantities, such as the total Casimir energy, are the same. Motivated by the possibility for the stabilization of extra dimensions by quantum effects, the one-loop Casimir energy of various braneworld compactifications has been investigated by several authors for both scalar and fermion fields (see references cited in Introduction). It has been shown that for the simultaneous solution of the stabilization and hierarchy problems a fine tuning of the model parameters is essential. From the point of view of backreaction of quantum effects the investigation of local densities is also of considerable interest. For a conformally coupled massless scalar field the corresponding results can be obtained from the results in the Minkowski spacetime by using the standard formula for conformally related problems. For the case of plane-symmetric line element (2.1) with an arbitrary function $\sigma(y)$ this has been done in Ref. [47] by using the results from [69]. Recently the energy-momentum tensor in the Randall-Sundrum braneworld for a bulk scalar with zero brane mass terms c_1 and c_2 is considered in Ref. [49]. This paper appeared when our calculations were in progress. Note that in [49] only a general formula is given for the unrenormalized VEV in terms of the differential operator acting on the Green function. In our approach the application of the generalized Abel-Plana formula allowed us to extract manifestly the part due to the AdS bulk without boundaries and for the points away from the boundaries the renormalization procedure is the same as for the boundary-free parts. The latter is well-investigated in literature. In addition, the boundary induced parts are presented in terms of exponentially convergent integrals convenient for numerical calculations.

6 Conclusion

The natural appearance of AdS in a variety of situations has stimulated considerable interest in the behavior of quantum fields propagating in this background. In the present paper we have investigated the Wightman function and the vacuum expectation value of the energy-momentum tensor for a scalar field with an arbitrary curvature coupling parameter satisfying Robin boundary conditions on two parallel plates in AdS spacetime. The application of the generalized Abel-Plana formula to the mode sum over zeros of the corresponding combinations

of the cylinder functions allowed us to extract the boundary-free AdS part and to present the boundary induced parts in terms of exponentially convergent integrals. In the region between two plates the Wightman function is presented in two equivalent forms given by Eqs. (2.31) and (2.43). The first terms on the right of these formulae are the Wightman function for AdS bulk without boundaries. The second ones are induced by a single plate and the third terms are due to the presence of the second plate. The expectation values for the energy-momentum tensor are obtained by applying on the Wightman function a certain second order differential operator and taking the coincidence limit. For the case of a single plate geometry this leads to formula (3.12) for the region $z > z_a$ and to formula (3.15) for the region $z < z_a$. As we could expect from the problem symmetry the part of this tensor corresponding to the components on the hyperplane parallel to the plate is proportional to the corresponding metric tensor. On the boundary the vacuum energy-momentum tensor diverges, except the case of a conformally coupled massless scalar. The leading terms of the corresponding asymptotic expansion near the boundary are given by expressions (3.26) and (3.27). These terms are the same as for a plate in the Minkowski bulk. They do not depend on the Robin coefficient, have different signs for Dirichlet and Neumann scalars and vanish for a conformally coupled scalar. The coefficients for the subleading asymptotic terms will depend on the AdS curvature radius, Robin coefficient and on the mass of the field. For large proper distances from the plate to compared with the AdS curvature radius, $k_D|y - a| \gg 1$, the boundary induced energy-momentum tensor vanishes as $\exp[2\nu k_D(a - y)]$ in the region $y > a$ and as $\exp[k_D(2\nu + D)(y - a)]$ in the region $y < a$. The same behavior takes place for a fixed $y - a$ and large values of the parameter k_D . In the large mass limit, $m \gg k_D$, the boundary parts are exponentially suppressed. Note that here we consider the bulk energy-momentum tensor. For a scalar field on manifolds with boundaries in addition to this part, the energy-momentum tensor contains a contribution located on the boundary (for the expression of the surface energy-momentum tensor in the case of arbitrary bulk and boundary geometries see Ref. [74]). As it has been discussed in Refs. [73, 69, 65, 75, 76, 74], the surface part of the energy-momentum tensor is essential in considerations of the relation between local and global characteristics in the Casimir effect. The vacuum expectation value of the surface energy-momentum tensor for the geometry of two parallel branes in AdS bulk is evaluated in Ref. [77]. It is shown that for large distances between the branes the induced surface densities give rise to an exponentially suppressed cosmological constant on the brane. In particular, in the Randall-Sundrum model the cosmological constant generated on the visible brane is of the right order of magnitude with the value suggested by the cosmological observations.

For the case of two-plates geometry the vacuum expectation value of the bulk energy-momentum tensor is presented as a sum of purely AdS, single plates, and 'interference' parts. The latter is given by Eq. (4.5) and is finite for all values $z_a \leq z \leq z_b$. In the limit $z_a \rightarrow z_b$ the standard result for two parallel plates in the Minkowski bulk is obtained. For two-plates case the vacuum forces acting on boundaries contain two terms. The first ones are the forces acting on a single boundary when the second boundary is absent. Due to the well-known surface divergencies in the expectation values of the energy-momentum tensor these forces are infinite and need an additional regularization. The another terms in the vacuum forces are finite and are induced by the presence of the second boundary and correspond to the interaction forces between the plates. These forces are given by formula (4.12) with $j = a, b$ for the plate at $z = z_a$ and $z = z_b$ respectively. For Dirichlet scalar they are always attractive. In the case of mixed boundary conditions the interaction forces can be either repulsive or attractive. Moreover, there is a region in the space of Robin parameters in which the interaction forces are repulsive for small distances and are attractive for large distances. This provides a possibility to stabilize interplate distance by using the vacuum forces. For large distances between the plates, the vacuum interaction forces per unit surface are exponentially suppressed by the factor $\exp[2\nu k_D(a - b)]$ for the plate at $y = a$ and by the factor $\exp[k_D(2\nu + D)(a - b)]$ for the plate at $y = b$. In Section

5 we give an application of our results for two plates case to the Randall-Sundrum braneworld with arbitrary mass terms on the branes. For the untwisted scalar the Robin coefficients are expressed through these mass terms and the curvature coupling parameter. For the twisted scalar Dirichlet boundary conditions are obtained on both branes. Note that in this paper we have considered boundary induced vacuum densities which are finite away from the boundaries. As it has been mentioned in Ref. [78], the same results will be obtained in the model where instead of externally imposed boundary condition the fluctuating field is coupled to a smooth background potential that implements the boundary condition in a certain limit [79].

Acknowledgments

I am grateful to Michael Bordag and Dmitri Vassilevich for valuable discussions and suggestions. The work was supported by the DAAD grant. I acknowledge the hospitality of the Institute for Theoretical Physics, University of Leipzig.

References

- [1] C. Fronsdal, Phys. Rev. D **10**, 589 (1974).
- [2] C. Fronsdal and R. B. Haugen, Phys. Rev. D **12**, 3810 (1975)
- [3] S. J. Avis, C. J. Isham, and D. Storey, Phys. Rev. D **18**, 3565 (1978).
- [4] P. Breitenlohner and D. Z. Freedman, Ann. Phys. (N.Y.) **144**, 249 (1982).
- [5] W. Heidenreich, Phys. Lett. **B110**, 461 (1982).
- [6] B. Binengar, C. Fronsdal, and W. Heidenreich, Ann. Phys. (N.Y.) **149**, 254 (1983); J. Math. Phys. **24**, 2828 (1983).
- [7] E. S. Fradkin and A. A. Tseytlin, Nucl. Phys. B **234**, 472 (1984).
- [8] N. Sakai and Y. Tanii, Phys. Lett. **B146**, 38 (1984).
- [9] C. Dullemond and E. van Beveren, J. Math. Phys. **26**, 2050 (1985).
- [10] C. P. Burgess and C. A. Lütken, Phys. Lett. **B153**, 137 (1985).
- [11] B. Allen and T. Jacobson, Commun. Math. Phys. **103**, 669 (1986).
- [12] R. Camporesi, Phys. Rev. D **43**, 3958 (1991).
- [13] R. Camporesi and A. Higuchi, Phys. Rev. D **45**, 3951 (1992).
- [14] M. Kamela and C. P. Burgess, Can. J. Phys. **77**, 85 (1999).
- [15] L.-X. Li, Phys. Rev. D **59**, 084016 (1999).
- [16] M. M. Caldarelli, Nucl. Phys. B **549**, 499 (1999).
- [17] W. D. Goldberger and I. Z. Rothstein, Phys. Rev. Lett. **89**, 131601 (2002); W. D. Goldberger and I. Z. Rothstein, hep-th/0208060.
- [18] O. Aharony, S. S. Gubser, J. Maldacena, H. Ooguri, and Y. Oz, Phys. Rep. **323**, 183 (2000).

- [19] N. Arkani-Hamed, S. Dimopoulos, and G. Dvali, Phys. Lett. **B429**, 263 (1998); Phys. Rev. D **59**, 086004 (1999); I. Antoniadis, N. Arkani-Hamed, S. Dimopoulos, and G. Dvali, Phys. Lett. **B436**, 257 (1998).
- [20] L. Randall and R. Sundrum, Phys. Rev. Lett. **83**, 3370 (1999).
- [21] L. Randall and R. Sundrum, Phys. Rev. Lett. **83**, 4690 (1999).
- [22] T. Gherghetta and A. Pomarol, Nucl. Phys. B **586**, 141 (2000).
- [23] A. Pomarol, Phys. Lett. **B486**, 153 (2000).
- [24] H. Davoudiasl, J. L. Hewett, and T. G. Rizzo, Phys. Lett. **B473**, 43 (2000).
- [25] R. Altendorfer, J. Bagger, and D. Nemeschansky, Phys. Rev. D **63**, 125025 (2001).
- [26] S. R. Huber and Q. Shafi, Phys. Rev. D **63**, 045010 (2001).
- [27] M. Gell-Mann, B. Zwiebach, Phys. Lett. B **141**, 333 (1984); Nucl. Phys. **B260**, 569 (1985).
- [28] W. D. Goldberger and M. B. Wise, Phys. Rev. Lett. **83**, 4922 (1999); Phys. Rev. D **60**, 107505 (1999); C. Csaki, J. Erlich, T. Hollowood, and Y. Shirman, Nucl. Phys. B **581**, 309 (2000); K. Maeda and D. Wands, Phys. Rev. D **62**, 124009 (2000); C. Barcelo and M. Visser, Phys. Rev. D **63**, 024004 (2001).
- [29] V. M. Mostepanenko and N. N. Trunov, *The Casimir Effect and Its Applications* (Clarendon, Oxford, 1997).
- [30] G. Plunien, B. Muller, and W. Greiner, Phys. Rep. **134**, 87 (1986).
- [31] *The Casimir Effect. 50 Years Later* edited by M. Bordag (World Scientific, Singapore, 1999).
- [32] M. Bordag, U. Mohideen, and V. M. Mostepanenko, Phys. Rep. **353**, 1 (2001).
- [33] M. Bordag, ed., Proceedings of the Fifth Workshop on Quantum Field Theory under the Influence of External Conditions, Int. J. Mod. Phys. A **17** (2002), No. 6&7.
- [34] K. A. Milton, *The Casimir Effect: Physical Manifestation of Zero-Point Energy* (World Scientific, Singapore, 2002).
- [35] M. Fabinger and P. Horava, Nucl. Phys. B **580**, 243 (2000).
- [36] S. Nojiri, S. Odintsov, and S. Zerbini, Phys. Rev. D **62**, 064006 (2000).
- [37] S. Nojiri and S. Odintsov, Phys. Lett. **B484**, 149 (2000).
- [38] D. J. Toms, Phys. Lett. **B484**, 119 (2000).
- [39] S. Nojiri, O. Obregon, and S. Odintsov, Phys. Rev. D **62**, 104003 (2000).
- [40] W. Goldberger and I. Rothstein, Phys. Lett. **B491**, 339 (2000).
- [41] S. Nojiri and S. Odintsov, JHEP **0007**, 049 (2000).
- [42] A. Flachi and D. J. Toms, Nucl. Phys. B **599**, 305 (2001).
- [43] A. Flachi and D. J. Toms, Nucl. Phys. B **610**, 144 (2001).

- [44] I. Brevik, K. A. Milton, S. Nojiri, and S. D. Odintsov, Nucl. Phys. B **599**, 305 (2001).
- [45] J. Garriga, O. Pujolas, and T. Tanaka, Nucl. Phys. B **605**, 192 (2001).
- [46] A. Flachi, I. G. Moss, and D. J. Toms, Phys. Lett. **B518**, 153 (2001); Phys. Rev. D **64**, 105029 (2001).
- [47] A. A. Saharian and M. R. Setare, Phys. Lett. **B552**, 119 (2003).
- [48] E. Elizalde, S. Nojiri, S. D. Odintsov, and S. Ogushi, Phys. Rev. D **67**, 063515 (2003).
- [49] A. Knapman and D. J. Toms, Phys. Rev. D **69**, 044023 (2004).
- [50] M. Bordag and D. V. Vassilevich, J. Phys. A **32**, 8247 (1999).
- [51] I. G. Moss, Phys. Lett. **B491**, 203 (2000).
- [52] P. B. Gilkey, K. Kirsten, and D. V. Vassilevich, Nucl. Phys. B **601**, 125 (2001).
- [53] S. Nojiri, S. Odintsov, and S. Zerbini, Class. Quantum Grav. **17**, 4855 (2000).
- [54] W. Naylor and M. Sasaki, Phys. Lett. **B542**, 289 (2002).
- [55] I. G. Moss, W. Naylor, W. Santiago-Germán, and M. Sasaki, Phys. Rev. D **67** 125010 (2003).
- [56] S. Mukohyama, Phys. Rev. D **63**, 044008 (2001).
- [57] R. Hofmann, P. Kanti, and M. Pospelov, Phys. Rev. D **63**, 124020 (2001).
- [58] A. H. Yeranyan and A. A. Saharian, Astrophysics **46**, 386 (2003).
- [59] J. Ambjorn and S. Wolfram, Ann. Phys. (N.Y.) **147**, 33 (1983).
- [60] H. Luckock, J. Math. Phys. **32**, 1755 (1991).
- [61] G. Esposito, A. Yu. Kamenshchik, and G. Polifrone, *Euclidean Quantum Gravity on Manifolds with Boundary* (Kluwer, Dordrecht, 1997).
- [62] S. N. Solodukhin, Phys. Rev. D **63**, 044002 (2001).
- [63] N. D. Birrell and P. C. W. Davis, *Quantum Fields in Curved Space* (Cambridge University Press, 1982).
- [64] A. A. Saharian, "The generalized Abel-Plana formula. Applications to Bessel functions and Casimir effect," Report No. IC/2000/14, hep-th/0002239.
- [65] A. A. Saharian, Phys. Rev. D **63**, 125007 (2001).
- [66] A. A. Saharian, Izv. AN Arm. SSR. Matematika **22**, 166 (1987) [Sov. J. Contemp. Math. Analysis **22**, 70 (1987)]; A. A. Saharian, Ph.D. thesis, Yerevan, 1987 (in Russian).
- [67] *Handbook of Mathematical Functions* edited by M. Abramowitz and I. A. Stegun (Dover, New York, 1972).
- [68] A. P. Prudnikov, Yu. A. Brychkov, and O. I. Marichev, *Integrals and series* (Gordon and Breach, New York, 1986), Vol.2.
- [69] A. Romeo, A. A. Saharian, J. Phys. A **35**, 1297 (2002).

- [70] P. Candelas and J. S. Dowker, Phys. Rev. D **19**, 2902 (1979).
- [71] H. H. Matevosyan and A. A. Saharian, to be published.
- [72] D. Deutsch and P. Candelas, Phys. Rev. D **20**, 3063 (1979).
- [73] G. Kennedy, R. Critchley, and J. S. Dowker, Ann. Phys. (N.Y.) **125**, 346 (1980).
- [74] A. A. Saharian, Phys. Rev. D **69**, 085005 (2004).
- [75] A. Romeo and A. A. Saharian, Phys. Rev. D **63** 105019 (2001).
- [76] S. A. Fulling, J. Phys. A **36**, 6857 (2003).
- [77] A. A. Saharian, Phys. Rev. D **70**, 064026 (2004).
- [78] N. Graham, R. L. Jaffe, V. Khemani, M. Quandt, M. Scandurra, and H. Weigel, Phys. Lett. **B572**, 196 (2003).
- [79] N. Graham, R. L. Jaffe, V. Khemani, M. Quandt, M. Scandurra, and H. Weigel, Nucl. Phys. B **645**, 49, (2002); N. Graham, R. L. Jaffe, and H. Weigel, Int. J. Mod. Phys. A **17**, 846 (2002).